

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/16, 14/47, A61K 38/17, 39/21	A1	(11) International Publication Number: WO 95/33768 (43) International Publication Date: 14 December 1995 (14.12.95)
(21) International Application Number: PCT/GB95/01285 (22) International Filing Date: 2 June 1995 (02.06.95) (30) Priority Data: 9411534.2 8 June 1994 (08.06.94) GB (71)(72) Applicant and Inventor: HOLMS, Rupert. Donald [GB/GB]: 66 Regent's Park Road, London NW1 7SX (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN. European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: AIDS PROPHYLACTICS		
(57) Abstract <p>It is an object of the present invention to provide preparations for the treatment or prophylaxis of AIDS and systemic lupus erythematosus and related disorders. The invention is based on my discovery of the process triggered in the immune system by HIV which leads to AIDS. HIV has a specific mechanism to activate the immune system to allow it to replicate and this same immune activation leads to an autoimmune process which eventually leads to AIDS. The preparation for treatment comprises of novel synthesised peptides whose amino acid sequences are derived from a human protein called ezrin and are based on the following sequence: NH₂ThrGluLysLysArgArgGluThrValGluArgGluLysGluCOOH. The mechanism of action of these peptides is that they switch off specific retrovirally induced immune responses by immunological tolerance.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Title:

AIDS PROPHYLACTICS

References Cited

Patent Documents filed in Great Britain

- 5 GB 9411534.2, Holms R.D., 8th June 1994

OTHER PUBLICATIONS

- Could specific oral tolerance be a therapy for autoimmune disease,
1 0 Stephen H., Thompson G., Staines N.A
Immunol. Today 1990 **11**: 396-399
- Induction of immunity and oral tolerance with polymorphic class II major
histocompatibility complex alloptides in the rat,
1 5 Sayegh M.H., Khoury S.J., Hancock W.W., Weiner H.L., Carpenter C.B.
Proc Natl Acad Sci USA 1992 **89**: 7762-7766
- Oral tolerance in experimental autoimmune encephalomyelitis,
Whitacre C C, Gienapp I.E., Orosz C.G., Bitar D.M.,
2 0 *The Journal of Immunology* 1991 **147** 2155-2163
- Prevention of experimental autoimmune myasthenia gravis by manipulation of the
immune network with a complementary peptide for the acetylcholine receptor,
Araga S., LeBoeuf R.D., Blalock J.E.
2 5 *Proc Natl Acad Sci* 1993 **90**: 8747-8751
- Peripheral T-cell tolerance induced in naive and primed mice by subcutaneous injection
of peptides from the major cat allergen Fel dI,
Briner T.J., Kuo M., Keating K.M., Rogers B.L., Greenstein J.L.,
3 0 *Proc Natl Acad Sci* 1993 **90** 7608-7612
- Identification of Ezrin as an 81-kDa Tyrosine-phosphorylated protein in T cells Egerton
M., Burgess W.H., Chen D., Druker B.J., Bretscher A., Samelson L.E. *The Journal of*
Immunology 1992 **149**: 1847-1852
3 5
- Identification of the Two Major Epidermal Growth factor-induced Tyrosine
Phosphorylation Sites in the Microvillar Core Protein Ezrin,
Krieg J., Hunter T. *Journal of Biological Chemistry* 1992 **267**: 19258-19265

BACKGROUND OF THE INVENTION

The field of the present invention relates to the prevention and treatment of Acquired Immune Deficiency Syndrome (AIDS), systemic lupus erythematosus (SLE) and related diseases. As used herein, the various titles, and other such headings are intended for guidance, and should not be construed as limiting on the present invention.

Over the last fifteen years a major international research effort has been directed at developing effective vaccines and therapies for AIDS but has so far only achieved very limited success. AIDS is a disease which results from chronic infection by Human Immunodeficiency Virus (HIV). At least ten million people worldwide and at least one million people in the USA are infected with HIV. It is widely accepted that AIDS is caused by HIV but the precisely how HIV infection leads to AIDS is still a matter of active scientific investigation. Current research into developing a vaccine to prevent AIDS is based on attempting to increase immune responses to the HIV but has so far failed to produce an effective product. There are now many thousands publications which describe attempts to develop AIDS therapies and vaccines by stimulating the human immune system to recognise and destroy the virus. The main problem appears to be that antibodies and cytotoxic T cell responses stimulated by prototype vaccines fail to protect against either HIV infection or AIDS disease progression because of the rapid mutation rate of HIV.

A number of drugs have been developed for the treatment of AIDS (mainly anti-viral drugs with a similar mode of action to 3'-azido-3'-deoxythymidine, AZT), but none have been demonstrated to prevent the development of AIDS after HIV infection. The main problem is that HIV mutates rapidly leading to changes in the virally encoded proteins which are the target for antiviral drugs and drug resistance develops quickly.

Systemic lupus erythematosus (SLE) is a disease which is normally viewed as being completely unrelated to AIDS. SLE is an autoimmune disease and effects mainly women. Over half a million women in the USA suffer from mild to severe SLE. SLE is an autoimmune disease with many immunological abnormalities such as lymphadenopathy, hypergammaglobulinemia, leukopenia, deposition of antigen-antibody complexes and autoantibodies resulting in chronic generalized connective tissue disorders ranging from mild to severe marked by skin eruptions, arthralgia, arthritis, anemia, visceral lesions, neurologic manifestations, fever and other constitution symptoms. Symptoms fluctuate in intensity over many years and SLE is normally controlled with immunosuppressive drugs ranging from non-steroidal anti-inflammatory drugs to immunosuppressive steroids but there is no effective treatment. SLE may be the result of immune activation

triggered by a retrovirus but no retrovirus has been identified which has been shown to cause SLE. GB 9411534.2 filed by Holms R.D, on 8th June 1994 discloses how HIV induces an autoimmune mechanism related to SLE which eventually leads to AIDS.

5 SUMMARY OF THE INVENTION

The present invention is based on selectively decreasing certain immune responses to HIV using peptides based on human proteins particularly a human protein called ezrin, providing a novel and non-obvious solution to the problem of the prevention and treatment of AIDS. A number of investigators have described autoimmunity related to HIV infection but have not established that HIV induced autoimmunity directly causes AIDS. The present invention is based on my discovery of the molecular mechanism of HIV induced autoimmunity which leads to AIDS and parallels between HIV induced disease and autoimmune disease such as systemic lupus erythematosus (SLE). The present invention provides for preparations which inhibit autoimmune processes which lead to AIDS and SLE by the induction certain types of immunosuppression or immune tolerance. GB 9411534.2 filed by Holms R.D, on 8th June 1994 discloses therapies and vaccines for the treatment and prevention of AIDS based on this novel autoimmune mechanism induced by HIV.

I propose that just before the HIV epidemic commenced, HIV acquired a small piece of human DNA which allowed it to evolve a specific mechanism to replicate more efficiently by activating the human immune system. Unfortunately, HIV encoded derivatives of this human sequence not only activate the system but also induce autoimmunity and apoptosis. When a peptide derived from the virus, in a complex with certain types of MHC, is seen as sufficiently foreign to induce an immune response but has sufficient sequence homology with human peptides to trigger T cells which recognise both viral and human antigens, there is a serious risk of a chronic autoimmune response. This type of molecular mimicry has been postulated for the induction of a number of autoimmune diseases including multiple sclerosis. Mammalian immune systems also tend to be strongly activated by foreign antigenic peptides which are very similar but not the same as self antigens.

The HIV amino acid sequence;
NH₂ThrLysAlaLysArgArgValValGluArgGluLysArgCOOH,
at position 498 to 510 (in the conserved C4 region) at the carboxy-terminus of HIV gp120 (a predicted alpha-helical region) is encoded between the direct repeats of a putative transposable element and appears to be a recent addition of human DNA to the virus. It is an immunodominant region of gp120 in man and accounts for up to 70% of

the total antibody response generated to the virus in some individuals although the antibodies are not virus-neutralising. A peptide derived from this region is also a dominant MHC Class-I restricted epitope for the induction of Cytotoxic T Lymphocytes (CTL). The sequence is conserved in all isolates of HIV-1 and HIV-2 against a background of a high mutation frequency (it is even more stable than the CD4 binding site) but it is different from an equivalent SIV sequence.

This 13 amino acid sequence of HIV; NH₂ThrLysAlaLysArgArgValValGluArgGluLysArgCOOH, is likely to be an epitope for the induction of autoimmunity as it has a high degree of homology (50% or greater) with certain human sequences. I have named amino acid sequences in HIV which mimic human sequences; Virus Homologous Peptide (referred to hereinafter as VHP) and the above HIV amino acid sequence VHP1. I have named the group of human amino acid sequences derived from endogenous human protein sequences which the virus mimics, Human Endogenous Peptide (referred to hereinafter as HEP).

Although I propose the following novel immunological process to describe the mechanism of action of these human peptides (HEPs), the following immunological process serves to illustrate the invention only, and should not be construed as limiting it in any way.

I suggest that there is a positive selection mechanism for mutants of HIV which maintain the HIV sequence VHP1 (NH₂ThrLysAlaLysArgArgValValGluArgGluLysArgCOOH) at position 498 to 510 (conserved C4 region) at the C-terminus of HIV gp120. This sequence is maintained because when it is presented by some MHC Class I molecules (particularly MHC Class I B8 and MHC Class I B35) and most MHC Class II DR molecules, it is important for inducing large numbers of activated T cells in which HIV infection and replication occurs. Unfortunately, the activated CD4+ T cells die or are killed rapidly after VHP1 activation and a population of VHP1 activated CD8+ T cells raise the level of autoreactive responses. Eventually the combination of both immunological processes leads to AIDS.

Chronic HIV infection results in continuous presentation of VHP1 produced by HIV and continuous stimulation of the immune system. Although the presence of activated T cells is essential for HIV replication, HIV only infects and kills a small minority of the VHP1 activated T cells. The majority of the activated T cells die as a result of VHP1 over stimulation or are destroyed by negative feedback mechanisms in the immune system involving autoreactive T cells. T cells are continually produced by the body to make good

these losses in a chronically infected HIV patient but the net result is a progressive decline in CD4+ T cells over a number of years.

I also propose that the continuous presentation of VHP peptide by certain MHC Class I molecules of HIV infected cells breaks the tolerance of the host immune system to self peptides such as those derived from ezrin and other HEPs presented on activated T and B cells. The general level of activation leads to the stimulation of autoreactive cytotoxic T cells and chronic B cell activation with the over production of a range of antibody molecules including autoreactive antibody. A key step in the activation of the autoimmune process in HIV infected individuals is the stimulation of autoreactive CD8+ cells which carry cross reactive T cell Receptor (TCR) molecules that recognise both VHP and HEP.

It is an object of this invention to prevent and treat AIDS and SLE and related disorders by the induction of specific immunological tolerance which inhibits the above pathological mechanism. It is an object of the invention to provide composition of matter comprising pharmaceutical grade purified peptide or a mixture of two or more different peptides up to thirty amino acids long or derivative molecules with additional chemical groups attached to such peptides, comprising an amino acid sequence with at least 50% homology over fourteen consecutive amino acids with the following human amino acid sequence:

NH₂ThrGluLysLysArgArgGluThrValGluArgGluLysGluCOOH,

(the fourteen amino acid sequence hereinafter referred to as HEP1) and a purified peptide or mixture of two or more different peptides up to thirty amino acids long or derivative molecules with additional chemical groups attached to such peptides, comprising an amino acid sequence of at least five consecutive amino acids with 100% homology with HEP1. It is a further object to provide a purified peptide comprising an amino acid sequence sufficiently duplicative of the fourteen amino acid sequence HEP1. It is a further object of the invention to provide a preparation of comprising peptides and peptide derivatives with at least 50% homology over fourteen consecutive amino acids with HEP1 and a purified peptide or mixture of two or more different peptides up to thirty amino acids long or derivative molecules with additional chemical groups attached to such peptides, comprising an amino acid sequence of at least five consecutive amino acids with 100% homology with HEP1, for the prevention and treatment of AIDS and related disorders which inhibits *in vivo* in man, partially or completely, HIV virus as measured by either an HIV p24 antigen assay *in vitro* or by an HIV infectivity assay *in vitro*. It is a further object of this invention to provide a preparation comprising peptides and peptide derivatives with at least 50% homology over fourteen consecutive amino acids with HEP1 and a purified peptide or mixture of two or more different peptides up to

thirty amino acids long or derivative molecules with additional chemical groups attached to such peptides, comprising an amino acid sequence of at least five consecutive amino acids with 100% homology with HEP1, for the prevention and treatment of AIDS and related disorders, and for the prevention and treatment of Systemic Lupus Erythematosus and related disorders, which inhibits *in vivo* in man, partially or completely, autoimmune or autoreactive responses measured *in vitro* in a T cell Proliferation Assay.

The close similarity between foreign VHP1 and self HEP1 when presented by certain MHC molecules and the possible immunoregulatory role of HEP1 in the healthy immune system, leads to strong stimulatory signals being transmitted through the immune system during HIV infection. The administration of HEP1 above the threshold for immunological tolerance induction is likely to block the VHP1 mediated activation of the immune system by HIV infected T cells. The clinical result of HEP1 administration predicted from the above model of HIV induced autoimmunity is lower levels of immune activation, a rise in CD4 T cell levels and a fall in the level of chronic HIV infection. (Figure 1)

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 entitled 'Immune activation and HIV infection' is a schematic diagram to show how an HIV infected CD4+ cell activates other CD4+ and CD8+ cells by the presentation of VHP on MHC molecules. The minority of CD4+ cells activated by this process are then infected by HIV while the majority of CD4+ T cells become over stimulated and die. Some of the activated CD8+ T cells are autoreactive and activate an autoimmune process which kills or suppresses activated T cells expressing HEP. The administration of HEP1 blocks the HIV induced T cell activation process and inhibits chronic HIV infection.

Figure 2 entitled 'Inhibition of HIV by Orally Administered HEP1 *in vivo* in patient PP' is a graphical representation of the normalised clinical trial data in Example 2 (data rebased to 100 on day 1 of the trial). The graph shows the result of oral administration of HEP1: the decline in HIV virus particles as measured by p24Ag levels, the decline in HIV infectivity as measured by a TCID assay and the fall in total white count (decline in immune activation).

Figure 3 entitled 'Improvement of Immune Status by Subcutaneously Administered HEP1 *in vivo* in patient PP' is a graphical representation of the normalised clinical trial data in Example 2 (data rebased to 100 on day 1 of the trial). The graph shows the result of subcutaneous administration of one larger dose of HEP1: a sharp increase in CD4+ and CD8+ T cells.

DETAILED DESCRIPTION OF THE INVENTION

5 Induction of immune tolerance with HEP1 related peptides and the prevention and treatment of AIDS

The non-obvious similarities between the HIV amino acid sequence; NH₂ThrLysAlaLysArgArgValValGluArgGluLysArgCOOH, position 498 to 510 (conserved C4 region) at the C-terminus of HIV gp120 (VHP1) and sequences in human proteins, were investigated using computer searches of the SwissProt protein sequences database (Release 25). I established that VHP1 has a 70% sequence homology to an evolutionarily conserved human protein called ezrin between amino-acid positions 324-337 of ezrin. The result of the search for homology is set forth in the table below.

	HIV gp120-gp41 Cleavage site-X													
Position	498	499	500	501	502	503	504	505	506	507	508	509	510	
gp120	Thr	Lys	Ala	Lys	Arg	Arg	Val	Val	Glu	Arg	Glu	Lys	Arg	
Homology														
Ezrin	Thr	Glu	Lys	Lys	Arg	Arg	Glu	Thr	Val	Glu	Arg	Glu	Lys	Glu
Position	324	325	326	327	328	329	330	331	332	333	334	335	336	337

Carboxy terminus of amino acid sequences- x

I made the unexpected discovery that nine out of fourteen amino acids were identical when the HIV amino acid sequence between position 498 and 510 of gp120 was compared to the human amino acid sequence between positions 324 and 337 of ezrin. It is not obvious from prior publications that this region ezrin is important for immune regulation.

The fourteen amino acid sequence of HEP1, NH₂ThrGluLysLysArgArgGluThrValGluArgGluLysGluCOOH, is identical to the amino acid sequence between positions 324 and 337 of human ezrin. No publication had disclosed HEP1 related peptides in man.

Ezrin, a human tubulin binding protein, is found in the cytoplasm of T cells and is phosphorylated by tyrosine kinase during T cell activation. Ezrin is also known as P81.

Cytovillin or Villin-2, is a protein of 585 amino acids. The homology extends over a predicted alpha-helical region of ezrin and is adjacent to the tyrosine 353 phosphorylation site. Ezrin is part of a family of ezrin related proteins, which includes ezrin, radixin, moesin and merlin. All these proteins are related in the region that shows similarity to gp120, but ezrin has the most significant homology with 9/13 identities. Although ezrin behaves as a soluble protein in T cells and appears to be diffusely cytoplasmic by immunofluorescence staining, it is believed that the critical population of ezrin is that which is associated with the submembraneous cortical cytoskeleton (Identification of Ezrin as an 81-kDa Tyrosine-phosphorylated protein in T cells Egerton M., Burgess W.H., Chen D., Druker B.J., Bretscher A., Samelson L.E. *The Journal of Immunology* 1992 **149**: 1847-1852). It may also be important for the autoimmune pathology of AIDS that ezrin is also found localized to microvillar actin microfilament cores in the brush border of the intestinal epithelium and also in neurons. In the gut epithelium ezrin is involved in changes in cell membrane morphology in response to stimulation (Identification of the Two Major Epidermal Growth factor-induced Tyrosine Phosphorylation Sites in the Microvillar Core Protein Ezrin, Krieg J., Hunter T. *Journal of Biological Chemistry* 1992 **267**: 19258-19265).

In addition to Ezrin, other human (self) proteins also possess a weaker primary sequence homology with VHP1: these include Heat shock protein 89, NK-TR protein, T complex protein, Adrenergic receptor Type 2, Creatine Kinase, Trypsin inhibitor, Aldehyde Dehydrogenase, Opioid Receptor (kappa), Glycine Receptor (Alpha-2 chain) and Ryanodine receptor. Mouse Histone H2B (very similar to human Histone H2B) also has homology.

It is an object of this invention that induction of tolerance can be used to switch-off immune activation and autoimmunity induced by HIV to prevent and treat HIV infection and AIDS. The present invention provides a vaccine against AIDS comprising a toleragenic dose (significantly larger than the threshold dose sufficient to stimulate an immune response) of pharmaceutical grade purified peptide or a mixture of two or more different peptides up to thirty amino acids long or derivative molecules with additional chemical groups attached to such peptides, comprising an amino acid sequence with at least 50% homology over fourteen consecutive amino acids with HEP1 and a purified peptide or mixture of two or more different peptides up to thirty amino acids long or derivative molecules with additional chemical groups attached to such peptides, comprising an amino acid sequence of at least five consecutive amino acids with 100% homology with HEP1 or any combination thereof, or peptides similar thereto. The preferred route of administration is oral, although it is equally possible to achieve tolerance by sub-cutaneous, intravenous or intramuscular administration by standard

methods. It is an aspect of this invention that a peptide vaccine based on HEP1 can be administered to prevent and treat AIDS in HIV infected people by the induction of tolerance to the VHP sequences. It is another aspect of this invention that a peptide vaccine based on HEP1 can be administered to prevent and treat SLE in people by the
5 induction of tolerance to other HEP sequences.

HOW TO MAKE

Peptides used for the toleragenic vaccine may be synthesized, for example, using a solid
10 phase method using either Boc or Fmoc chemistry or any other practical route for peptide synthesis known to those skilled in the art of peptide synthesis. After synthesis, peptides are cleaved from the resin (in solid phase methods) and deprotected using trifluoromethane sulphonic acid or Hydrogen Fluoride or other agents. Peptides may be
15 desalted by column chromatography and purified by HPLC. Purity of the peptides may be demonstrated by reverse-phase HPLC or by automatic sequencing. Without limiting the possible formulations of the vaccine or therapy, the preferred composition comprises peptide or peptides either dissolved in sterile pharmaceutical saline or distilled water to a final concentration between 1 and 1000 mg/ml. The preparation may be freshly made up for each administration or may be stored frozen for up to 30 days at -20°C. (see
20 **EXAMPLE 1 & EXAMPLE 2**).

HOW TO USE

The induction of immunological tolerance by oral administration or sub cutaneous
25 administration of peptide solutions has been successfully achieved in various animal experiments. The immunological tolerance induced by peptides can occur either by induction of T cell anergy (direct inhibition) or by induction of suppressor T cell populations (indirect inhibition) by the peptides. Those skilled in the art of induction of immunological tolerance are aware of procedures to induce tolerance in animals by oral,
30 sub-cutaneous, intravenous or intramuscular administration of small peptides derived from cross reactive antigens. Examples of these procedures have been disclosed in the following publications:

Could specific oral tolerance be a therapy for autoimmune disease,
35 Stephen H., Thompson G., Staines N.A
Immunol. Today 1990 **11**: 396-399

Induction of immunity and oral tolerance with polymorphic class II major histocompatibility complex allopeptides in the rat.

Sayegh M.H., Khoury S.J., Hancock W.W., Weiner H.L., Carpenter C.B.
Proc Natl Acad Sci USA 1992 **89**: 7762-7766

- Oral tolerance in experimental autoimmune encephalomyelitis,
5 Whitacre C C, Gienapp I.E., Orosz C.G., Bitar D.M.,
The Journal of Immunology 1991 **147** 2155-2163

- Prevention of experimental autoimmune myasthenia gravis by manipulation of the
immune network with a complementary peptide for the acetylcholine receptor,
10 Araga S., LeBoeuf R.D., Blalock J.E.
Proc Natl Acad Sci 1993 **90**: 8747-8751

- Preipheral T-cell tolerance induced in naive and primed mice by subcutaneous injection
of peptides from the major cat allergen Fel dI,
15 Briner T.J., Kuo M., Keating K.M., Rogers B.L., Greenstein J.L.,
Proc Natl Acad Sci 1993 **90** 7608-7612

- Two factors are important for the induction of tolerance: high dose of soluble peptide
antigen and absence of any co-stimulatory particles or adjuvant. Generally,
20 immunological activation occurs with very much lower doses of peptide antigen than the
threshold dose above which the peptide antigen induces tolerance or inhibition of
immune responses (for example in man a very small dose of 10 nanograms of a small
foreign peptide antigen may illicit a strong immune response whereas a 10 milligram
dose may inhibit the immune response). The dose range for the induction of tolerance
25 varies between peptides but the amount or concentration may be determined
experimentally. Subcutaneous administration of peptides yield final serum concentrations
of peptide between 50-100X higher than by the oral route and therefore tolerance may be
achieved with less peptide than by the oral route. The peptide solution should be filter
sterilised to remove particulates and any microbiological contaminants, if a peptide
30 solution is to be administered by injection.

- It is an object of this invention that the preferred doses are between 1mg and 5000mg of
pharmaceutical grade purified peptide or a mixture of two or more different peptides up to
thirty amino acids long or derivative molecules with additional chemical groups attached
35 to such peptides, comprising an amino acid sequence with at least 50% homology over
fourteen consecutive amino acids with HEP1 and a purified peptide or mixture of two or
more different peptides up to thirty amino acids long or derivative molecules with
additional chemical groups attached to such peptides, comprising an amino acid sequence

of at least five consecutive amino acids with 100% homology with HEP1 or any combination thereof, or peptides similar thereto.

5 Although it is theoretically possible to attempt to induce immunological tolerance with peptides based on any HIV amino acid sequence, there is the general risk that a protective immune response against HIV may be inhibited. There is also a specific risk associated with administering peptides based on VHP1 to induce immunological tolerance as an incorrect dose of VHP1 based peptides may aggravate the disease by increasing immune activation. It is an advantage of this invention that the administration of peptides based on
10 human amino acid sequences are likely to be significantly safer than peptides based on HIV amino acid sequences.

The following Examples serve to illustrate the invention only, and should not be construed as limiting it in any way.

15

EXAMPLE 1

IN VITRO MODEL OF VHP1-INDUCED AUTOIMMUNITY IN HUMAN CELLS FROM UNINFECTED DONORS.

20 MATERIALS and METHODS

Peptide synthesis

VHP1 peptide and control peptide used in this study were synthesized using a solid phase method using Fmoc chemistry. They were cleaved from the resin and deprotected
25 using trifluoromethane sulphonic acid. Peptides were desalted on a P-10 column in PBS (pH 7.3). Homogeneity of the peptides was indicated by reverse-phase HPLC. Peptides were sequenced by automatic Edman degradation using gas phase sequence and were shown to be >95% pure. The following peptides were synthesised:

30 13 amino acid sequence of VHP1 derived from HIV:
NH₂ThrLysAlaLysArgArgValValGluArgGluLysArgCOOH

A 14 amino acid control sequence:

NH₂LeuGluAspArgArgAlaAlaValAspThrValCysArgAlaCOOH

35

Donors

All donors who volunteered for this study were HIV negative. The MHC phenotype of donors was determined before the experiment. The number of the subtype of the A, B

and C categories of MHC Class I phenotype and the number of the subtype of the DR, DP and DQ categories of MHC Class II phenotype of each donor is set out in the following tables:

5 Donors positive for B8 or B35 (*Trigger+*)

Donor	A	B	C	DR	DP	DQ
FM	1,3	8	7	3	nd	nd
SW	1,3	35,62	nd	1	nd	1
WW	11,36	35,61	nd	1	nd	1
JMC	1,2	8,45	6,7	3,4	nd	2,7
Target 1	1	8	7	3	3	2
Target 2	31	35	4	4	nd	7

Donors negative for B8 or B35 (*Trigger-*)

Donor	A	B	C	DR	DP	DQ
TM	3,28	7,14	nd	6	nd	nd
JF	2	17	3	6	nd	1
KB	1	7,14	7	4,11	nd	7
ML	2	44,52	5	2	nd	1
Target 3	3	27	1	1	nd	5

Generation of VHP1 induced short-term cultures

10

Peripheral Blood Mononuclear Cells (PBMC) were isolated from 8 HIV seronegative individuals who volunteered as donors for the study. Four donors FM, SW, WW, JMC (*Trigger+*) were positive for either MHC B35 or B8 and four donors TM, JF, KB, ML (*Trigger-*) were negative for MHC B35 or B8. PBMC ($10^6/\text{ml}$) from the each of the

15

donors were incubated in the presence of VHP1 or control peptide at 100 ng/ml for 4-8 hours to produce Effector Cells. Irradiated autologous PBMC without VHP1 stimulation served as controls.

Cytotoxicity Assay

20

VHP1 induced short-term cultures of autoreactive T cells (Effector Cells) were assayed for cytotoxic activity against autologous PHA T cell blasts stimulated with 5 $\mu\text{g}/\text{ml}$ PHA for 4-5 days at a concentration of 10^4 cells/well in the absence of either control or VHP1 peptide (Target Cells). $1-2 \times 10^6$ Target Cells were labeled with 250mCi of ^{51}Cr in 0.3

25

ml of medium for 1 hour at 37°C. Target cells were washed 3x in medium, diluted to 10^5 or 10^4 cells/ml and 100ml of cell suspension added to each well of a V-bottomed 96-well

microtitre plate. Effector Cells were added to the Target Cells at Effector:Target (E:T) ratios from 50:1 to 6.25:1. Cultures were incubated for 4 hours at 37°C and supernatants were harvested and counted for ⁵¹Cr release. Percent cytotoxicity is calculated using the formula : 100x (experimental release - spontaneous release)/(maximum release-spontaneous release). Spontaneous release and maximum release were determined by incubating target cells in medium or 1.0% Triton x-100 respectively. Results were calculated as means ± Standard Deviations (SD) of triplicate cultures and a positive result was scored above 20% specific lysis. The cytotoxicity assay was repeated for allogeneic Target cells from donors (Target-1, Target-2 and Target-3) and the percent specific lysis scored at an Effector:Target ratio of 10:1.

T cell Proliferation Assay

The proliferation assays was performed by culturing either PBMC or T cell lines (10⁴ cells/well) in U-bottomed 96-well microtitre plates for 3 days at 37°C stimulated with fresh irradiated autologous PBMC and r-IL2 either with or without 1μM VHP1. PBMC or T cell lines were rested in the absence of r-IL2 for 24 hours prior to assay. Three positive control cultures of the PBMC or T cell lines were stimulated with 10μg/ml PHA, 10μg/ml PPD and 0.1μg/ml SEB (Staph. Enterotoxin) respectively to determine maximum proliferation. The two negative controls were an unstimulated culture of PBMC or T cell line and irradiated PBMC or T cell line under the same conditions. During the final 6 hours before harvesting, the cultures received 1mCi/well of [³H] thymidine. The cells were harvested onto glass fibre filters by a semiautomatic harvester and the incorporation of labeled thymidine was quantitated by liquid scintillation spectrometry. The results are expressed as counts per minute (cpm) of the mean for triplicate cultures.

RESULTS

Induction of autoreactive cells by VHP1

Short term cultures of PBMC from donors positive for B8 or B35 (*trigger+*) stimulated with VHP1 produced Effectors which showed significant autocytotoxic responses against activated autologous T cell targets in the Cr release cytotoxicity assays. Short term cultures of PBMC from donors positive for B8 or B35 stimulated with control peptide (Control) and short term cultures of PBMC from donors negative for B8 or B35 (*trigger-*) did not produce significant levels these autocytotoxic Effectors. The dependence of autocytotoxic responses on MHC phenotype of uninfected donors in short term cultures of PBMC stimulated with VHP1 is shown below.

Dependence of autoreactive response to VHP1 on MHC Phenotype
of Donor

	Peptide stimulant	
	VHP1	Control
DONOR		
Trigger+		
FM	+	-
SW	+	-
WW	+	-
JMC	+	-
Trigger-		
TM	-	-
JF	-	-
KB	-	-
ML	-	-

5 Identification of the target of VHP1 Induced Autoreactive Cells

To establish the MHC specificity of the cytotoxic cells stimulated by VHP1, further Cytotoxicity Assays with allogeneic PHA activated T cell Targets with different MHC phenotypes were then performed. The Effectors were derived from short term cultures of PBMC from donors of either MHC B8 or B35 (*trigger+*) incubated with VHP1 for 4-8 hours. The Target Cells were from donors with either B8 B35 (*trigger+*) or B27 (*Trigger-*) MHC phenotype. Target PBMC incubated with radioactive ^{51}Cr for one hour before being mixed with Effectors. Effector Cells were mixed with radiolabelled Target Cells and incubated for 4 hours and lysis was measured by release of radioactive ^{51}Cr .

15

EFFECTORS

	Donor	MHC Phenotype	Predicted Sensitivity to AIDS
	JMC	B8	High
	WW	B35	High
20	SW	B35	High

TARGETS

	Donor	MHC Phenotype	Predicted Sensitivity to AIDS
	1	B8	High
	2	B35	High
25	3	B27	Low

The measurement of percent specific lysis by ^{51}Cr release at an Effector to Target Ratio of 10:1.

5	Targets (Donor-MHC)	Effectors (Donor-MHC)		
		JMC-B8	WW-B35	SW-B35
	Target1-B8	30	30	9
	Target2-B35	30	32	10
	Target3-B27	7	14	2

10

JMC-B8 Effectors efficiently hit both B8 and B35 Targets but not B27 Targets. WW-B35 Effectors efficiently hit both B8 and B35 Targets but not B27 Targets (the same specificity as the JMC-B8 Effectors). SW-B35 Effectors showed a similar pattern of specificity in that they recognised both B8 and B35 Targets but not B27 Targets but the

15

overall cytotoxic activity from this donor was lower.

20

B8 and B35 both have the identical trigger amino acid sequence in the alpha-one domain of MHC Class I whereas B27 has a different sequence.(GB 9411534.2 filed by Holms R.D, on 8th June 1994). B8 and B35 phenotypes have been associated with high sensitivity to the development of AIDS after HIV infection in a number of clinical studies. I conclude that VHP1 is presented by MHC Class I B8 and B35 to Effector T cells in the short term cultures of PBMCs. I also conclude that the TCR of B8 and B35 Effector T cells are recognising a common epitope on MHC Class I B8 and B35 of the Targets which is absent on MHC Class I B27. The common epitope is likely to be a combination of the common MHC Class I B trigger sequence (present in the alpha one domain of both B8 and B35 but not on B27) plus human peptides resembling HEP1 (self peptides homologous to VHP1).

25

30

An autoreactive T cell line was established from donor WW (HLA-B35 phenotype) by stimulation with VHP1. The line was a mixed population of 59% CD4+T cells and 28% CD8+ T cells. A proliferation assay was performed to measure to level of proliferation in the T cell line to autologous (self) cells. T cell line WW responded vigorously to autoantigens presented on autologous irradiated PBMC at similar levels to the PHA or PPD stimulated positive control cultures.

Discussion

I conclude that VHP1 (at low concentrations) activates the immune system and induces significant levels of autolytotoxic cells in donors with MHC phenotypes associated with sensitivity to developing AIDS rapidly (B8 or B35) while it is not so active in donors who do not carry the MHC phenotypes associated with sensitivity to rapid development of AIDS. The correlation between VHP1 sensitivity and sensitivity to rapid development of AIDS suggests that the expression and correct presentation of the VHP1 sequence in HIV infected people may be an important step in the disease process which leads to AIDS.

EXAMPLE 2

CLINICAL TRIAL OF HEP1 IN AN HIV INFECTED PATIENT

The objective of this study was to determine in an HIV infected patient if the induction of immunological tolerance to HEP1 could reduce immune activation and HIV infection. The design of the first clinical experiment was to orally administer HEP1 to the patient in daily 10mg doses (the lower end of the preferred dose range). This was then followed approximately three months later with a subcutaneous administration of HEP1 at a higher dose of 140mg. To assess the therapeutic benefit of HEP1, the following disease progression markers were measured by taking blood samples before, during and after each trial and determining: HIV levels by a p24 antigen assay, HIV infectivity by a cellular TCID assay and immune activation by quantitating levels of various lymphocytes.

2.5 The Patient

Patient PP volunteered to take peptide HEP1 orally while attending the HIV Clinic at Ealing Hospital NHS Trust. Patient PP is a white male, 32 years old and has been HIV infected for eight years and entered the trial with a T cell count of 80 cells / mm³. He had not taken any anti-viral therapy (for example AZT) in the previous five years before the trial. At the time of the trial, the patient was HCV negative (Abbott 2nd generation), HepB negative and Blood group A rhesus (D) positive.

Synthesis of HEP1 peptide

3.5 Abbreviations

DCC - Dicyclohexylcarbodiimide

DIC - Diisopropylcarbodiimide

DCM - Dichloromethane

DMF - Dimethylformamide

TFA - Trifluoroacetic acid

Boc - t-Butyloxycarbonyl-

HOBT - Hydroxybenzotriazole

DIEA - Diisopropylethylamine

5 DCU - Dicyclohexylurea

HF - Hydrogen Fluoride

370mg of HEP1 (a 14 amino acid peptide) was synthesized by Boc-synthesis based on a human ezrin sequence:

1 0 $\text{NH}_2\text{ThrGluLysLysArgArgGluThrValGluArgGluLysGluCOOH}$.

0.05 mmole resin (Boc-aminoacid -OCH₂ pam resin) and a three fold excess of activated Boc-aminoacid solution (activated with 0.5M HBTU in DMF and 2.5 mmole DIEA) were coupled in the following steps. The resin was washed with DMF, Boc protecting groups

1 5 minute, drained, activated amino acid solution was added, shaken for 10 minutes at room temperature then washed and the sequence of steps repeated for the next amino acid. On

completion of the synthesis, the reaction mixture was flow washed with DMF, then with DCM and dried. The peptide was then cleaved from the resin with HF (-5°C for 1.5

2 0 hours), the HF was evaporated, the mixture was washed with 50ml ether and evaporated then the peptide dissolved in 6M guanidine 0.1M TRIS for HPLC preparative separation. After separation, the peptide was analysed for purity by HPLC and was determined to be over 99% pure HEP1. The peptide solution was then evaporated to a pure white fluffy solid under vacuum.

2 5 **Preparation of HEP1 peptide for Oral Administration.**

To a vial containing 221mg of pure freeze dried HEP1 peptide approximately 5ml of sterile distilled water was added to dissolve the peptide and the solution was washed into a 50 ml volumetric flask and made up to exactly 50 ml in the volumetric flask with more

3 0 sterile distilled water giving a final concentration of 4.42 mg/ml HEP1. 2.26 ml volumes of the solution were transferred into twenty two separate lots in 5 ml plastic tubes and stored at -20°C (10mg of peptide per tube). Two vials were stored at -70°C as reference solutions at the clinic and the remaining 20 vials were provided to the patient on dry ice in a vacuum flask. The vials were stored until use by the patient in a freezer at -20°C. The

3 5 patient thawed out one tube (leaving the remaining tubes frozen in the Freezer) each morning and swallowed the solution one hour before breakfast. The procedure was repeated for twenty days.

Preparation of HEP1 peptide for Subcutaneous Administration.

HEP1 was administered subcutaneously to patient PP three months after the end of the oral administration study.

5

Preparation

To a vial containing 147 mg of pure freeze dried HEP1 peptide (no inorganic salts added) 1ml of sterile distilled water was added to the vial to dissolve the peptide giving a
1 0 final concentration of 147 mg/ml HEP1. The solution was filter sterilised through a 0.2 micrometer sartorius filter attached to a steril 5 ml syringe and then transferred a 5ml sterile container.

Cellular TCID Assay

1 5

Cellular TCID is a technique that measures the relative amount of viral load within cells. Within four hours of a sample being taken from the HIV infected patient, 10^6 viable PBMC from the sample were obtained by centrifugation of the anticoagulated blood over Ficoll-Hypaque lymphocyte separation medium. A serial dilution of the patients cells was
2 0 set up in a 24 well plate in decreasing amounts (1:1, 1:4, 1:16, 1:64 and 1:256 and a negative control) using 0.9 ml special media (RPMI-1640) supplemented with 15% fetal calf serum and 10% interleukin 2. Infected cells were then cocultivated with 10^6 PBMC derived from an uninfected human donor that had been stimulated for 24-48 hours with phytohaemagglutinin (PHA) to activate the cells to make them susceptible to infection by
2 5 HIV. The cocultures were subsequently monitored for the presence of p24 antigen in the supernatant fluid twice weekly for up to 14 days, during this period the plates were spun on days 4,7,11 and 14 and then a 50% medium change was performed. The medium was removed at days 7 and 14 and saved in duplicate 0.5ml eppendorff tubes (minimum 250 μ l/tube) for future p24 antigen testing. The medium was removed on day 11 and any
3 0 remaining medium from day 14 (after saving for p24 antigen testing) was pooled and stored in liquid nitrogen for future virus culture. A culture was considered positive if the concentration of p24 antigen in the supernatant on day 7 and day 14 exceeded 30 picogram/ml (typical cut-off value). The lowest number of peripheral mononuclear cells required to produce a positive culture was taken as the end point and the reciprocal of the
3 5 end point dilution indicated the relative number of infected cells in the patient.

p24Ag Assay

p24Ag is a marker of HIV infection as well as a predictor of HIV disease progression. The p24Ag assay measures p24 antigen from the core of HIV and is an immunoassay using murine monoclonal antibody coated onto microwell strips. The assay detects HIV in plasma, serum or tissue culture medium. If present, the antigen binds to the antibody-coated microwells. The bound antigen is recognised by biotinylated antibodies to HIV which react with conjugated streptavidin-horseradish peroxidase. Colour develops from the reaction of the peroxidase with hydrogen peroxide in the presence of tetramethylbenzidine (TMB) substrate. Specimens which are HIV p24 Ag positive can be quantitated by setting up a standard curve using serial dilutions of the Antigen Reagent. The assay is not accurate below 30 picograms per ml.

CD4, CD8, NK

T cell subsets were measured by adding a whole blood sample to a 4.5ml tube containing EDTA and staining with monoclonal antibodies to CD3, CD4 and CD8. The percentage of T cells in Peripheral Blood Lymphocytes were determined by flow-cytometric procedures. The numbers of CD4 and CD8 T cells were determined by obtaining total and differential white-cell counts and multiplying by the appropriate factor obtained on flow cytometry.

Results: Oral Administration of HEP1

Blood samples were taken from the patient 69 days before the trial and on the first day of the trial before the first dose of HEP1, to determine the background levels of the disease progression markers. Oral administration of HEP1 solution containing 10mg HEP1 was started on day 2 of the trial and continued until day 22. Further Blood samples were taken for analysis on day 7, day 14, day 21 and day 28 of the trial (7 days after the last dose of HEP1). The patient experienced no adverse reactions and continued to feel well throughout the administration of HEP1.

The following markers of HIV disease progression were assayed: p24 Ag (the concentration of HIV p24 protein in picogram/ml in the blood), TCID (the infectivity of the HIV in the cells of blood sample measured by an *in vitro* cell culture assay in which a serial dilution of the sample was mixed with activated uninfected PBMC and the level of HIV infection determined by p24Ag assay), total white count (immune activation measured by the number of white cells per mm^3), CD4 (the number of CD4 positive cells per mm^3), CD8 (the number of CD8 positive cells per mm^3), NK (the number of Natural

Killer cells per mm^3), Lymphocytes (the number of cells per mm^3), Monocytes (the number of cells per mm^3), and Granulocytes (the number of cells per mm^3). Normalised data was calculated by dividing each data point by the value of its data series on day 1 of the trail and multiplying by 100.

5

	BEFORE TREATMENT		DURING TREATMENT			AFTER
Trial day	day -69	day 1	day 7	day 14	day 21	day 28
Date	3/11/94	10/1/95	17/1/95	24/1/95	1/2/95	7/2/95
p24Ag	n/a	71	44	n/a	n/a	38
TCID	n/a	1:16	n/a	n/a	n/a	1:1
White count	7900	8200	8200	7700	6800	5500
CD4	90	80	80	90	80	80
CD8	850	900	850	900	910	730
NK	20	30	30	40	40	20
Lymphocytes	1260	1480	1150	1460	1160	1050
Monocytes	630	570	490	620	540	440
Granulocytes	6000	6230	6560	5620	5100	4020
%Lymphocytes	15.4	17.8	15.5	16.8	18.6	20
%Monocytes	4.2	6.1	3.1	6.4	5.2	5.5
%Granulocytes	80.4	76.1	81.4	76.8	76.2	74.5

NORMALISED DATA- day 1 = 100

Trial day	day -69	day 1	day 7	day 14	day 21	day 28
Date	3/11/94	10/1/95	17/1/95	24/1/95	1/2/95	7/2/95
p24Ag		100	62			54
TCID		100				6
White count	96	100	100	94	83	67
CD4	113	100	100	113	100	100
CD8	94	100	94	100	101	81
NK	67	100	100	133	133	67
Lymphocytes	85	100	78	99	78	71
Monocytes	111	100	86	109	95	77
Granulocytes	96	10	105	90	82	65
%Lymphocytes	87	100	87	94	104	112
%Monocytes	96	100	51	105	85	90
%Granulocytes	106	100	107	101	100	98

These preliminary clinical results show that orally administered HEP1 reduced viral load by 46% as measured by p24Ag, and reduced the infectivity of HIV by 16X as measured by TCID and that the effect persisted for 7 days after treatment. It is important to note that the p24Ag level was only just above background at 38 picogram/ml as 30 picogram/ml is considered the typical cut-off value. The total white cell count was rising before treatment but fell during and after treatment suggesting a reduction in immune activation had been achieved during treatment. The persistence of these effects for seven days after treatment suggests that a T suppressor cell population had been induced in the patient which reduced infectivity of HIV and immune activation. CD8 increased during treatment but there was no significant increase in CD4 count (at this concentration of HEP1). (Figure 2)

Results

1.5 Subcutaneous Administration of HEP1

0.95 ml (the volume equivalent to 140mg HEP1) of the steril solution was self-injected subcutaneously in the abdominal wall in one dose on day 1 of the subcutaneous HEP1 trial. Patient PP experienced a stinging sensation which lasted approximately 15 minutes but there were no other adverse reactions. Blood samples were taken on day -92, day -57, day -8, day 1 before the HEP1 injection and day 1* four hours after the injection and then 6 days and 13 days after the HEP1 injection. The samples were assayed for p24, TCID, CD4, CD8, total white count, lymphocytes, granulocytes, monocytes and the data is set out in the table below (normalised data was calculated as described above).

2.5

	End Oral study	BEFORE TREATMENT				AFTER
Trial day	day -91	day -56	day -7	day 1	day 1*	day 6
Date	7/2/95	14/3/95	2/5/95	9/5/95	10/5/95	16/5/95
p24Ag	38	n/a	n/a	n/a	n/a	n/a
TCID	1:1	n/a	n/a	n/a	n/a	n/a
White count	5500	6600	8200	5600	4600	5400
CD4	80	75	70	80	120	80
CD8	730	880	670	880	1130	780
NK	20	75	40	30	30	
Lymphocytes	1050	1250	980	1290	1520	
Monocytes	440	400	570	450	510	
Granulocytes	4020	4950	6640	3860	2530	
%Lymphocytes	20	19	12.9	23.3	33	22.2
%Monocytes	5.5	6	7	8	11	
%Granulocytes	74.5	75	81	69	55	

NORMALISED DATA- day 1 = 100

Trial day	day -91	day -56	day -7	day 1	day 1*	day 6
Date	7/2/95					
p24Ag	n/m	n/a	n/a	n/a	n/a	n/a
TCID	n/m	n/a	n/a	n/a	n/a	n/a
White count	98	118	146	100	82	96
CD4	100	94	88	100	150	100
CD8	83	100	76	100	128	89
NK	67	250	133	100	267	
Lymphocytes	81	97	76	100	118	
Monocytes	98	89	127	100	113	
Granulocytes	104	128	172	100	66	
%Lymphocytes	86	82	55	100	142	
%Monocytes	69	75	88	100	138	
%Granulocytes	108	109	117	100	80	

- 5 The subcutaneous administration of one larger dose of HEP1 resulted in a sharp rise in both CD4 T cells (+50%) and CD8 T cells (+28%) after the HEP1 injection followed by a return to the pre-injection levels six days later (p24Ag and TCID data was not ready in time for the filing of the application). (Figure 3)

AIDS PROPHYLACTICS

CLAIMS

5

WHAT IS CLAIMED IS:

- 10 1. Composition of matter comprising a purified peptide or mixture
of two or more different peptides up to thirty amino acids long or
derivative molecules with additional chemical groups attached to such
peptides, comprising an amino acid sequence with at least 50% homology
over fourteen consecutive amino acids with the following amino acid
15 sequence:
NH₂ThrGluLysLysArgArgGluThrValGluArgGluLysGluCOOH,
(the fourteen amino acid sequence hereinafter referred to as HEP1) and a
purified peptide or mixture of two or more different peptides up to thirty
amino acids long or derivative molecules with additional chemical groups
20 attached to such peptides, comprising an amino acid sequence of at least
five consecutive amino acids with 100% homology with HEP1.
2. A purified peptide of claim 1 comprising an amino acid sequence
sufficiently duplicative of the fourteen amino acid sequence HEP1.
- 25 3. A preparation of claim 1 for the prevention and treatment of AIDS
and related disorders which inhibits *in vivo* in man, partially or
completely, HIV virus as measured by either an HIV p24 antigen assay *in*
vitro or by an HIV infectivity assay *in vitro*.
- 30 4. A preparation of claim 1 for the prevention and treatment of AIDS
and related disorders, and for the prevention and treatment of Systemic
Lupus Erythematosus and related disorders, which inhibits *in vivo* in
man, partially or completely, autoimmune or autoreactive responses
35 measured *in vitro* in a T cell Proliferation Assay.

Figure 1
IMMUNE ACTIVATION AND HIV INFECTION




Figure 2
Inhibition of HIV
by orally administered HEP1
in vivo in Patient PP

Normalised data: Day1 = 100






Figure 3
Improvement of immune status
by subcutaneously administered HEP1
in vivo in Patient PP

Normalised data: Day1 = 100



INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/GB 95/01285

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K14/16 C07K14/47 A61K38/17 A61K39/21

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 05196 (HIVER LTD) 2 April 1992 see page 63 - page 66; claims 1,2,14-21; examples 7,10 ---	1-4
X	WO,A,92 22572 (REPLICO MEDICAL AB) 23 December 1992 see claims 1,8,9; tables 1A-1C ---	1-4
A	WO,A,94 03487 (ZAGURY JEAN FRANCOIS) 17 February 1994 see the whole document ---	1-4
A	WO,A,93 15750 (HAYNES BARTON F) 19 August 1993 see the whole document ---	1-4
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

27 September 1995

Date of mailing of the international search report

10. 10. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Groenendijk, M

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/GB 95/01285

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO,A,92 00755 (HOFFMANN GEOFFREY W ;KION TRACY A (CA); WELDER CLAYTON A (CA)) 23 January 1992 see the whole document -----</p>	1-4

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/01285

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9205196	02-04-92	AU-B- 641814 AU-A- 8653691 EP-A- 0502160	30-09-93 15-04-92 09-09-92
WO-A-9222572	23-12-92	AU-A- 2023092	12-01-93
WO-A-9403487	17-02-94	FR-A- 2694560 FR-A- 2694938 AU-B- 4712993 EP-A- 0656010	11-02-94 25-02-94 03-03-94 07-06-95
WO-A-9315750	19-08-93	AU-B- 3662993 EP-A- 0652764 JP-T- 7506810	03-09-93 17-05-95 27-07-95
WO-A-9200755	23-01-92	US-A- 5230887 AU-A- 8180191 EP-A- 0589875	27-07-93 04-02-92 06-04-94